

Modelling of growth cycle of water hyacinth: An application to Bolgoda Lake

By

Miss. P. U. Dilani Fernando

LIBRARY
UNIVERSITY OF MORATUWA
CIVIL ENGINEERING DEPARTMENT

THIS THESIS WAS SUBMITTED TO THE DEPARTMENT OF
CIVIL ENGINEERING OF THE UNIVERSITY OF MORATUWA
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR
THE DEGREE OF MASTER OF SCIENCE IN ENVIRONMENTAL
WATER RESOURCES ENGINEERING AND MANAGEMENT



6/2/03
5.5.12.201

MSc. in
Water Resources Eng
at night
UM Thesis coll

80524

Department of Civil Engineering, University of Moratuwa, Sri Lanka

March 2003

University of Moratuwa




80524

80524


DECLARATION

I hereby declare that the work included in the thesis in part or whole has not been submitted in any form for any other academic qualification of any institution.



Miss. P. U. Dilani Fernando

Certified by



Dr. Mahesh Jayaweera

Supervisor



University of Moratuwa, Sri Lanka.
Electronic Theses & Dissertations
www.lib.mrt.ac.lk

ABSTRACT

Water hyacinth (*Eichhornia crassipes* (Mart). Solms) is considered as a problematic aquatic weed in many lakes, irrigation canals, stagnant ponds, waterways and semi-wet areas in Sri Lanka. Bolgoda Lake is one of the freshwater bodies in Sri Lanka, which has been severely affected by excessive growth of water hyacinth thereby clogging the water ways and hence adversely affecting water quality. This study was conducted to determine the growth characteristic of water hyacinth under influence of natural, physical and chemical factors in Bolgoda Lake. The parameters considered in the study were as follows: biomass in dry weight, biomass production per day, phosphorus and nitrogen contents in plant tissues, phosphorus and nitrogen contents in the water body, pH, temperature, and salinity.

The luxuriant growth of water hyacinth was observed during the study period, which occurred with the temperature ranging between 26-32 °C, pH from 6.67-7.76, salinity from 0-1.5 ppt and water nutrients from 4.6-17.4 mg N/l, 0.18-0.70 mg NO₃-N/l and 0.14-0.93 mg P/l and 0.02-0.16 mg PO₄-P/l respectively. Under such conditions, results revealed that hyacinth plants produced a biomass yield of 20 –1800 g dry weight/m² and the number of plants increased from 21 to 412 per m² for the entire study period of 14 weeks with doubling time of around 13-15 days. The biomass production rate varied from 2.10-75.25 g dry weight/m² per day. Results of heavy metal uptake experiment suggest that rhizofiltration (metal absorption into roots) and phytoextraction (concentrate into the harvestable parts of roots or shoots) are the key mechanisms for removal of heavy metals from the aqueous phase. Phytoextraction was more responsible in translocating heavy metals to aboveground parts in initial few weeks and rhizofiltration became prominent at the later stages in which more metals are bound to below-ground parts. Once the heavy metal binding was complete, harvesting was suggested at the end of the 13th week during which more metals were adsorbed only to root zone.

From this study it was shown that there exists a massive proliferation of water hyacinth stands in Bolgoda Lake with a great influence of nitrogen, phosphorus, pH and temperature. However, there has been a significant perishment of the existing

stands of the vegetation from time to time due to the exposure to saline waters entering from the tidal action.

A numerical model was developed to simulate the growth of water hyacinth in Bolgoda Lake, Sri Lanka. The model was first applied to experimental data from Sato and Kondo, (1981). Secondly, it was used to evaluate the management options to control the growth of water hyacinth in Bolgoda Lake. Model application showed how the model could be used to evaluates the management options to control the growth of water hyacinth and to reduce the available nutrients in the system. These options include harvest strategies (initial density and harvesting interval) and harvest rate. The maximum yield of 329 g / m^2 dry weight was obtained when the rate of harvest was analogous to the initial density (at 100 g dry wt/m^2) in that the water hyacinths were harvested at a uniform rate every 20 days. The continuous harvesting is the major objective criteria to remove available nutrients in the water body and to control the excessive growth of water hyacinth in Bolgoda Lake.



ACKNOWLEDGEMENT

I would like to express my deep gratitude to my project supervisors, Dr. Nimal Priyantha Gamage and Dr. Mahesh Jayaweera, Senior Lecturers, Department of Civil Engineering, University of Moratuwa, Sri Lanka for their help and encouragement throughout my study. Their support and guidance have proven invaluable both personally and professionally. Also I profoundly extend my sincere appreciation to Dr. S. L. J. Wijekoon and Dr. S. A. S. Kulathilaka, Senior lecturers of University of Moratuwa, Sri Lanka for their support throughout the study.

I gratefully acknowledge the financial support granted by the Department of Civil Engineering, University of Moratuwa, Sri Lanka. I also wish to acknowledge Professor Takashi Asaeda, Saitama University, Japan, for providing me with chemicals necessary for experimental work.

I am thankful to Professor (Mrs.) N. Rathnayake, Director Postgraduate studies, and Dr. G. Kodikara, Former Head of the Department of Civil Engineering, University of Moratuwa, Sri Lanka for giving me this valuable opportunity and granting permission to carry out the research at University premises.

Also I would like to express my gratitude to Miss. Priyanka Dissanayake, Analytical Chemist, Mrs. Nilanthi Gunathilaka, Technical Officer and Mr. L. Justin Silva, Lab. attendant in Environmental Engineering Laboratory, University of Moratuwa, Sri Lanka for their kind support and assistance given to me throughout the study period.

At last but not least I sincerely wish to express my deepest gratitude to my mother, and friends for encouraging me in pursuing the research work.

TABLE OF CONTENTS

	<u>Page No.</u>
List of Tables	i
List of Figures	ii
List of Abbreviations	iv
List of Symbols	vii
<u>Chapter One: Introduction</u>	1
1.0 General introduction	2
1.1 Brief description of water hyacinth	3
1.2 Generalization life cycle of <i>Eichhornia crassipes</i>	4
1.3 Appearance and spread of water hyacinth	5
1.4 Water hyacinth in Sri Lanka	6
1.5 Need for mathematical relationships and models	6
<u>Chapter Two: Literature Review</u>	8
2.0 Literature review	9
2.1 Influence of temperature, pH and salinity	10
2.2 Nitrogen availability and uptake	11
2.3 Phosphorus availability and uptake	12
2.4 Heavy metal availability and uptake	12
2.5 Growth and dry matter production	13
2.6 Modelling of growth of water hyacinth	14

<u>Chapter Three: Description of Study Site</u>	15
3.0 Description of study site	16
<u>Chapter Four: Methodology of Study</u>	19
4.0 Materials and methodology	20
4.1 Estimation of length of the shoot and the roots	20
4.2 Estimation of Biomass	21
4.3 Chemical analysis of plant materials	21
4.3.1 Analysis of total nitrogen and nitrate-nitrogen in plant tissues	21
4.3.2 Analysis of total phosphorus and reactive phosphate-phosphorus in plant tissues	22
4.4.3 Analysis of heavy metals in plant tissues	23
4.4 Chemical analysis of water	23
<u>Chapter Five: Model Formulation</u>	24
5.0 The mathematical equations in the model	25
5.1 State equations for growth kinetics	25
5.2 Respiration kinetics	29
5.3 Mortality kinetics	29
<u>Chapter Six: Results and Discussion</u>	30
6.0 Growth of plants and biomass	31
6.1 Number of plants	34
6.2 Influence of nutrients on growth of water hyacinth	35

6.2.1 Influence of total nitrogen and nitrate-N ($\text{NO}_3\text{-N}$)	36
on growth of water hyacinth	
6.2.2 Influence of total phosphorus and reactive phosphates-P	37
($\text{PO}_4\text{-P}$) on growth of water hyacinth	
6.3 Influence of salinity on biomass production	41
6.4 Influence of temperature on biomass production	44
6.5 Influence of pH on biomass production	45
6.6 Heavy metal uptake and its influence on growth of water hyacinth	46
6.6.1 Influence of chromium on growth	46
6.6.2 Influence of cadmium on growth	48
6.6.3 Influence of copper on growth	49
6.6.4 Influence of lead on growth	50
6.6.5 Influence of cobalt on growth	50
6.6.6 Influence of nickel on growth	51
6.7 Model calibration	55
6.8 Model application	57
<u>Chapter Seven: Conclusion and recommendations</u>	60
7.0 Conclusion of the present work and recommendation for future	61
work	
Bibliography	64

LIST OF TABLES

	<u>Page No.</u>
Table 1: Methods of extraction, analysis and quantification with references	22
Table 2: Maximum growth rate and biomass of water hyacinth reported in literature	33
Table 3: Maximum growth rate and biomass of water hyacinth reported in literature	35
Table 4: Nutrient composition in water hyacinth	39



University of Moratuwa, Sri Lanka.
Electronic Theses & Dissertations
www.lib.mrt.ac.lk

LIST OF FIGURES

	<u>Page No.</u>
Figure 1: Life cycle of water hyacinth	5
Figure 2: Location map of Bolgoda Lake showing the study site	17
Figure 3: Exclusive growth of water hyacinth in Bolgoda Lake	18
Figure 4: Artificial enclosure in Bolgoda Lake	20
Figure 5: Variation of water hyacinth biomass in the study area	31
Figure 6: Water hyacinth growth rate	32
Figure 7: Increase in plant number	34
Figure 8: Variation of total Nitrogen in different plant tissues and ambient water	36
Figure 9: Variation of total Nitrate-N in different plant tissues and ambient water	37
Figure 10: Variation of total phosphorus in different plant tissues and ambient water	38
Figure 11: Variation of Phosphate-P in different plant tissues and ambient water	38
Figure 12: Relation between salinity and biomass	42
Figure 13: Relation between salinity and growth rate	42
Figure 14: Variation of water temperature and air temperature	44
Figure 15: Relation between water temperature and growth rate	44
Figure 16: Relation between biomass and pH	46
Figure 17: Relation between pH and growth rate	46
Figure 18: Cr variation in different parts of water hyacinth	47

Figure 19: Relation between Cr level in water and biomass of roots	48
Figure 20: Cd variation in different parts of water hyacinth (<i>Eichhornia crassipes</i>)	48
Figure 21: Cu variation in different parts of water hyacinth (<i>Eichhornia crassipes</i>)	49
Figure 22: Pb variation in different parts of water hyacinth (<i>Eichhornia crassipes</i>)	50
Figure 23: Co variation in different parts of water hyacinth (<i>Eichhornia crassipes</i>)	51
Figure 24: Ni variation in different parts of water hyacinth (<i>Eichhornia crassipes</i>)	52
Figure 25: Relation between Ni level in water and root tissues	52
Figure 26: Calibration of water hyacinth growth model using measured data from 1-fold culture solution of Sato and Kondo (1981)	56
Figure 27: Calibration of water hyacinth growth model using measured data from 3-fold culture solution of Sato and Kondo (1981)	56
Figure 28: Simulation of water hyacinth growth model using measured data from Bolgoda Lake, Sri Lanka	57
Figure 29: Simulation of growth of water hyacinth after 30 days and 40 days of harvesting	58
Figure 30: Simulation of growth after harvesting to a dry weight of 50 and 100 g/m ² at 20 days intervals	58

LIST OF ABBREVIATIONS

Abbreviations

Name

Nitrate-N	Nitrate Nitrogen
Phosphate-P	Phosphate Phosphorus
Dw	Dry weight
Ww	Wet weight
ppt	Parts per thousand
T	Temperature
Cd	Cadmium
Cr	Chromium
Co	Cobalt
Cu	Coppers
Pb	Lead
Ni	Nickel
Nitrate-N	Nitrate Nitrogen
Phosphate-P	Phosphate Phosphorus
$b_{shi}(i)$	Dry weight of above-ground biomass at i th layer
B_{rt}	Root biomass
$Pg_{shi}(i)$	Gross growth rate of shoots at the i th layer
R_{shi}	Respiration rate of shoots
R_{rt}	Respiration rate of roots
D_{shi}	Mortality rate of shoots



University of Moratuwa, Sri Lanka
Electronic Theses & Dissertations
www.lib.mrt.ac.lk

Abbreviations

Name

D_{rt}	Mortality rate of roots
G_{rt}	Supply of photosynthesized material for root growth
H_{sh}	Harvesting rates of shoots
H_{rt}	Harvesting rates of roots
B_{sh}	Total shoot biomass
N	Number of layers in the plant canopy
Pg_{mx}	Maximal photosynthetic growth rate
$f(T)$	Dependency of growth on the temperature
$f(I)$	Response of shoots of water hyacinth to light intensity as a function
$f(N)$	Nitrogen limitation
$f(P)$	Phosphorus limitation
I_{PAR}	Photosynthetically active radiation
k_c	Light extinction coefficient within the canopy
K_{PAR}	Half saturation irradiance
F_i	Cumulative leaf area index of the stand above the layer I
N	Concentration of inorganic nitrogen
P	Concentration of phosphors in water
K_N	Half saturation constant for nitrogen



University of Moratuwa
Electronic Theses & Dissertations
www.lib.mrt.ac.lk

Abbreviations

Name

	uptake
K_p	Half saturation constant for phosphors
	Uptake
g_m	Maximum specific growth rate of roots
K_{rt}	Half saturation coefficient of root age
Age	Age of roots
β_{sh}	Rate coefficients for respiration of shoots
β_{rt}	Rate coefficients for respiration of roots
$^{\circ}C$	Centigrade
mt	metric tons
HCL	Hydrochloric acid



University of Moratuwa
Electronic Theses & Dissertations
www.lib.mrt.ac.lk

LIST OF SYMBOLS

Symbol

Name

θ

Arrhenius constant



University of Moratuwa, Sri Lanka.
Electronic Theses & Dissertations
www.lib.mrt.ac.lk